

## anti-human CD25 APC-conjugated

The antibody MEM-181 reacts with CD25 (Interleukin-2 receptor  $\alpha$  chain), a 55 kDa type I transmembrane glycoprotein expressed on activated B and T lymphocytes, activated monocytes/macrophages and on CD4<sup>+</sup> T lymphocytes (T regulatory cells); it is lost on resting B and T lymphocytes.

**CD25** (IL2R $\alpha$ , Tac) is a ligand-binding alpha subunit of interleukin 2 receptor (IL2R). Together with beta and gamma subunit CD25 constitutes the high affinity IL2R, whereas CD25 alone serves as the low affinity IL2R. CD25 expression rapidly increases upon T cell activation. The 55 kDa CD25 molecule is enzymatically cleaved and shed from the cell surface as a soluble 45 kDa s-Tac, whose concentration in serum can be used as a marker of T cell activation. Expression of CD25 indicates the neoplastic phenotype of mast cells. Humanized anti CD25 antibodies represent a useful tool to reduce the incidence of allograft rejection as well as the severity of graft versus host reaction, and radioimmunoconjugates of anti-CD25 antibodies can be used against CD25 expressing lymphomas.

<b>Clone:</b>	MEM-181
<b>Isotype:</b>	Mouse IgM
<b>Physical state:</b>	Liquid for direct use
<b>Form:</b>	purified antibody conjugated with cross-linked Allophycocyanin (APC)
<b>Buffer:</b>	PBS containing 1% BSA and 0.1% sodium azide (pH 7.4)
<b>Storage:</b>	Store at 4 °C. Do not freeze. Avoid prolonged exposure to light.
<b>Application:</b>	Flow Cytometry
<b>Reference:</b>	Leukocyte Typing VI. Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997)
<b>Quantity:</b>	1.0 ml
<b>Order N°:</b>	H12159A

**Warning:** Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32).

This material is offered for **research only**. Not for use in human. For in vitro use only. EuroBioSciences will not be held responsible for patent infringement or other violations that may occur with the use of our products.