

anti-human CD2 APC conjugated

The antibody LT2 reacts with CD2, a 50 kDa glycoprotein present on the human peripheral blood T lymphocytes and NK cells; also expressed by all thymocytes.

HLDA VI; WS Code T 6T-008

CD2 belongs to T lymphocyte glycoproteins of immunoglobulin superfamily. Its interaction with CD58 stabilizes adhesion between T cells and antigen presenting or target cells. Relatively low affinity of CD2 to CD58 (as measured in solution) is compensated within the two-dimensional cell-cell interface to provide tight adhesion. Moreover T cell activation induces increased CD2 expression and its lateral mobility, making easier contact between CD2 and CD58. Subsequently T cell activation causes fixation of CD58-CD2 at sites of cell-cell contact, thereby strengthening intercellular adhesion. CD2 deficiency reduces intestinal inflammation and helps to control infection.

Clone:	LT2
Isotype:	Mouse IgG2a
Physical state:	Liquid for direct use
Form:	purified antibody conjugated with cross-linked Allophycocyanin (APC)
Buffer:	PBS containing 1% BSA and 0.1% sodium azide (pH 7.4)
Storage:	Store at 4 °C. Do not freeze. Avoid prolonged exposure to light.
Application:	Flow Cytometry
Reference:	Leukocyte Typing VI. Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997)
Quantity:	1.0 ml
Order N°:	H12393A

Warning: Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32).

This material is offered for **research only**. Not for use in human. For in vitro use only. EuroBioSciences will not be held responsible for patent infringement or other violations that may occur with the use of our products.