

anti-human CD62L FITC-conjugated

The antibody LT-TD180 reacts with CD62L (L-selectin), a 74-95 kDa single chain type I glycoprotein expressed on most peripheral blood B lymphocytes, T lymphocytes, monocytes and granulocytes; it is also present on a subset of NK cells and certain hematopoietic malignant cells.

CD62L (L-selectin) is an adhesion glycoprotein that is constitutively expressed on the cell surface of leukocytes and mediates their homing to inflammatory sites and peripheral lymph nodes by enabling rolling along the venular wall. CD62L is also involved in activation-induced neutrophil aggregation. Activation-dependent CD62L shedding, however, counteracts neutrophil rolling. CD62L has also signaling roles including enhance of chemokine receptor expression. Similarly to CD62P, the major ligand of CD62L is PSGL-1 (P-selectin glycoprotein ligand-1).

Clone:	LT-TD180
Isotype:	Mouse IgG1
Physical state:	Liquid for direct use
Form:	purified antibody conjugated with Fluorescein isothiocyanate (FITC)
Buffer:	PBS containing 1% BSA and 0.1% sodium azide (pH 7.4)
Storage:	Store at 4 °C. Do not freeze. Avoid prolonged exposure to light.
Application:	Flow Cytometry
Reference:	von Andrian UH and others: 1993 Jul 1;82(1):182-91. Simon SI and others: 1995 Aug 1;155(3):1502-14. Ding Z and others: 2003 Jun 1;101(11):4245-52
Quantity:	1.0 ml
Order N°:	H12201F

Warning: Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32).

This material is offered for **research only**. Not for use in human. For in vitro use only. EuroBioSciences will not be held responsible for patent infringement or other violations that may occur with the use of our products.