

## anti-human CD80 FITC-conjugated

The antibody MEM-233 reacts with CD80 (B7-1), a 60 kDa single chain type I glycoprotein of immunoglobulin supergene family, expressed on professional antigen-presenting cells, such as dendritic cells, macrophages or activated B lymphocytes.

**CD80** (B7-1) and **CD86** (B7-2) are ligands of T cell critical costimulatory molecule **CD28** and of an inhibitory receptor **CTLA-4** (CD152). The both B7 molecules are expressed on professional antigen-presenting cells and are essential for T cell activation, the both molecules can also substitute for each other in this process. The question what are the differences in CD80 and CD86 competency has not been fully elucidated yet; there are still conflicts in results about their respective roles in initiation or sustaining of the T cell immune response.

<b>Clone:</b>	MEM-233
<b>Isotype:</b>	Mouse IgG1
<b>Physical state:</b>	Liquid for direct use
<b>Form:</b>	purified antibody conjugated with Fluorescein isothiocyanate (FITC)
<b>Buffer:</b>	PBS containing 1% BSA and 0.1% sodium azide (pH 7.4)
<b>Storage:</b>	Store at 4 °C. Do not freeze. Avoid prolonged exposure to light.
<b>Application:</b>	Flow Cytometry
<b>Reference:</b>	Vasilevko V and others: 2002 Mar;21(3):137-49. Yadav D and others: 2004 Sep 15;173(6):3631-9. Thomas IJ and others: 2007 Nov 1;179(9):5936-46
<b>Quantity:</b>	1.0 ml
<b>Order N°:</b>	H12208F

**Warning:** Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32).

This material is offered for **research only**. Not for use in human. For in vitro use only. EuroBioSciences will not be held responsible for patent infringement or other violations that may occur with the use of our products.