

anti-human CD37 FITC-conjugated

This antibody recognizes the CD37 antigen on human B cells. On flow cytometry it stains approximately 2-10% of human peripheral blood mononuclear cells strongly. This antibody has been studied at the 3. International Workshop on Human Leucocyte Differentiation Antigens.

CD1a, together with CD1b and c, belongs to group 1 of CD1 glycoproteins. These proteins serve as antigen-presenting molecules for a subset of T cells that responds to specific lipids and glycolipids found in the cell walls of bacterial pathogens or self-glycolipid antigens such as gangliosides, and they have also roles in antiviral immunity. Unlike CD1b, CD1a is excluded from late endosomal compartments and instead traffics independently in the recycling pathway of the early endocytic system, and CD1a antigen presentation is independent upon vesicular acidification.

Clone:	HH1
Isotype:	Mouse IgG1
Physical state:	Liquid for direct use
Form:	purified antibody conjugated with Fluorescein isothiocyanate (FITC)
Buffer:	PBS containing 1% BSA and 0.1% sodium azide (pH 7.4)
Storage:	Store at 4 °C. Do not freeze. Avoid prolonged exposure to light.
Application:	Flow Cytometry
Reference:	McMichael, A.J. et al. (eds.), Leucocyte typing 3., Oxford University Press (1987)
Quantity:	1.0 ml
Order N°:	H12417F

Warning: Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32).

This material is offered for **research only**. Not for use in human. For in vitro use only. EuroBioSciences will not be held responsible for patent infringement or other violations that may occur with the use of our products.