

## anti-human CD235a FITC-conjugated

The monoclonal antibody AME-1 is directed against the alpha-chain of glycophorin (molecular mass 43 kDa). In direct agglutination tests this monoclonal antibody reacts with normoblasts and erythrocytes. No reaction occurs with En(a-) erythrocytes and other normal peripheral blood cells. The monoclonal antibody is very useful for the recognition of malignant erythroblastic diseases.

**CD235a** (Glycophorin A, GPA) is a transmembrane sialoglycoprotein expressed on erythrocytes and their precursors. Similarly to glycophorin B (GPB), these molecules provide the cells with a large mucin-like surface, which minimalizes aggregation between erythrocytes in the circulation. GPA is the carrier of blood group M and N specificities, while GPB accounts for S, s and U specificities. CD235a is a receptor of Hsa, an Streptococcus adhesin.

<b>Clone:</b>	AME-1
<b>Isotype:</b>	Mouse IgG1
<b>Physical state:</b>	Liquid for direct use
<b>Form:</b>	purified antibody conjugated with Fluorescein isothiocyanate (FITC)
<b>Buffer:</b>	PBS containing 1% BSA and 0.1% sodium azide (pH 7.4)
<b>Storage:</b>	Store at 4 °C. Do not freeze. Avoid prolonged exposure to light.
<b>Application:</b>	Flow Cytometry
<b>Reference:</b>	Leukocyte Typing VII. and others: Oxford University Press (2002)
<b>Quantity:</b>	1.0 ml
<b>Order N°:</b>	H12460F

**Warning:** Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32).

This material is offered for **research only**. Not for use in human. For in vitro use only. EuroBioSciences will not be held responsible for patent infringement or other violations that may occur with the use of our products.