

anti-human CD11b FITC-conjugated

The antibody HI11b recognizes CD11b antigen (Mac-1), a 165 kDa type I transmembrane protein mainly expressed on monocytes, granulocytes and NK-cells. The CD11b mediates neutrophil and monocyte interactions with stimulated endothelium.

HLDA VI; WS Code BP 310; HLDA VI; WS Code M 18.

CD11b (integrin α M subunit) is a 165 kDa type I transmembrane glycoprotein that non-covalently associates with integrin β 2 subunit (CD18); expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/CD18 integrin (Mac-1, CR3) is highly expressed on NK cells, neutrophils, monocytes and less on macrophages. CD11b/CD18 integrin is implicated in various adhesive interactions of monocytes, macrophages and granulocytes, facilitating their diapedesis, as well as it mediates the uptake of complement coated particles, serving as a receptor for the iC3b fragment of the third complement component.

Clone:	HI11b
Isotype:	Mouse IgG1
Physical state:	Liquid for direct use
Form:	purified antibody conjugated with Fluorescein isothiocyanate (FITC)
Buffer:	PBS containing 1 % BSA and 0.09 % sodium azide (pH 7.2)
Storage:	Store at 4 °C. Do not freeze. Avoid prolonged exposure to light.
Application:	Flow Cytometry
Reference:	Leukocyte Typing VI. Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).
Quantity:	1.0 ml
Order N°:	H12520F

Warning: Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32).

This material is offered for **research only**. Not for use in human. For in vitro use only. EuroBioSciences will not be held responsible for patent infringement or other violations that may occur with the use of our products.