

anti-human CD44

The antibody MEM-85 reacts with both cell surface-expressed and soluble form of CD44 antigen (Phagocyte glycoprotein 1), a 80-95 kDa transmembrane glycoprotein (hyaladherin family) present on the most of cells and tissues (leukocytes, endothelial cells, mesenchymal cells, etc.); it is negative on platelets and hepatocytes.

CD44 is a transmembrane glycoprotein expressed on the surface of most cells, which serves as a receptor for hyaluronan. CD44 mediates angiogenesis, cell adhesion, proliferation and migration, it is thus important for lymphocyte activation, recirculation and homing, it can thus serve e.g. as a modulator of macrophage recruitment in response to pathogen. Although CD44 functions are essential for physiological activities of normal cells, elevated CD44 expression correlates with poor prognosis in many carcinomas, facilitating tumour growth and metastasis, antiapoptosis and directional motility of cancer cells.

Clone:	MEM-85
Isotype:	Mouse IgG2b
Physical state:	Purified from ascites by protein-G affinity chromatography
Purity:	> 95% (by SDS-PAGE)
Buffer:	PBS with 15 mM sodium azide (pH 7.4)
Storage Instruction:	Store at 4 °C. For long-term storage aliquot and store at -20°C. Avoid freeze/thaw cycles.
Application:	Flow Cytometry, Immunoprecipitation, Western Blotting, ELISA
Reference:	Leukocyte Typing VI., Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997)
Quantity:	0.1 mg
Order N°:	H12177

Warning: Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32).

This material is offered for **research only**. Not for use in human. For in vitro use only. EuroBioSciences will not be held responsible for patent infringement or other violations that may occur with the use of our products.