

anti-human CD58

The antibody MEM-63 reacts with CD58 (LFA-3), a 40-70 kDa glycoprotein distributed over many tissues, leukocytes, erythrocytes, endothelial cells, epithelial cells and fibroblasts. The antibody inhibits CD2/CD58 interactions.

CD58 (LFA-3) is an immunoglobulin family adhesion molecule expressed by both hematopoietic and non-hematopoietic cells (often on antigen presenting cells) and serving as ligand of CD2. This interaction is important for T cell-mediated immunity. CD58 is expressed in transmembrane form and in GPI-anchored form; the later is constitutively associated with protein kinases whereas the transmembrane form activates kinase activity upon triggering. CD58 is a powerful tool for detection of minimal residual disease in acute lymphocytic leukemia, and for evaluation of liver damage related with hepatitis B.

Clone:	MEM-63
Isotype:	Mouse IgG1
Species Reactivity:	Human, Porcine
Physical state:	Purified from ascites by protein-A affinity chromatography
Purity:	> 95% (by SDS-PAGE)
Buffer:	PBS with 15 mM sodium azide (pH 7.4)
Storage Instruction:	Store at 4 °C. For long-term storage aliquot and store at -20°C. Avoid freeze/thaw cycles.
Application:	Flow Cytometry, Immunoprecipitation, Western Blotting (non reducing conditions)
Reference:	Veltroni M and other: 2003 Nov;88(11):1245-52. Ariel O and other: 2004 Jun;16(6):667-73. Sheng L an dother: 2006 Jul 14;12(26):4237-40.
Quantity:	0.1 mg
Order N°:	H12197

Warning: Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32).

This material is offered for **research only**. Not for use in human. For in vitro use only. EuroBioSciences will not be held responsible for patent infringement or other violations that may occur with the use of our products.